

## **Historic, archived document**


Do not assume content reflects current scientific knowledge, policies, or practices.



# THE PLANT DISEASE REPORTER

Issued By

Division of Mycology and Disease Survey



Supplement 89

Crop Losses from Plant Diseases in the United States

1934

December 15, 1935



BUREAU OF PLANT INDUSTRY

UNITED STATES DEPARTMENT OF AGRICULTURE

U.S.D.A.

Crop Loss from Pest Diseases in the  
United States - 1941



BUREAU OF PLANT INDUSTRY

UNITED STATES DEPARTMENT OF AGRICULTURE

Supplementary to

Crop Loss from Pest Diseases in the United States - 1940

1941

December 18, 1941



January 29

Distribution of 1940 crop pest losses in the

ESTIMATES OF CROP LOSSES FROM DISEASES IN THE UNITED STATES - 1934

Compiled by

Jessie I. Wood

Plant Disease Reporter  
Supplement 89

December 15, 1935

CONTENTS

Foreword . . . . .	2
Graphic Summary of Losses, 1925-34 . . . . .	3-13
Wheat . . . . .	3
Oats. . . . .	4-5
Corn. . . . .	5-6
Peach . . . . .	13
Potato . . . . .	7-8
Sweet Potato . . . . .	9-10
Apple. . . . .	11-12
Tabular Summary of Losses, 1934 . . . . .	14-45
Wheat. . . . .	14
Barley . . . . .	16
Rye. . . . .	18
Oats . . . . .	19
Field Corn . . . . .	20
Sweet Corn . . . . .	22
Sweet Potato . . . . .	23
Potato . . . . .	24
Tomato . . . . .	26
Snap Bean. . . . .	30
Dry Bean . . . . .	33
Green Peas . . . . .	34
Cotton . . . . .	36
Sugar Beet . . . . .	37
Apple. . . . .	38
Pear . . . . .	40
Peach . . . . .	41
Cherry . . . . .	42
Grape. . . . .	43
Strawberry . . . . .	44

FOREWORD

These estimates of crop losses for the year 1934 have been computed and are presented in the same way as in previous years (Plant Dis. Reptr. Suppl. 83. 1932).

In addition to the tables showing reduction in yield for 1934, a graphic summary of losses during the ten years from 1925 to 1934 is furnished. This summary does not include all the crops or diseases for which loss estimates are given. The choice was made on the basis of a sufficient number of States reporting for a sufficient number of years to make a comparison possible. Some diseases showed so little variation from year to year that no graphs were prepared, although adequate estimates were available.

Besides the yearly fluctuation in loss, the graphs show the averages for the two overlapping ten-year periods 1920 to 1929 and 1925 to 1934. The earlier average is the higher in most cases, although sometimes so little higher that there is practically no difference. Exceptions are oat smuts, corn smut and ear rot, potato tipburn and leaf roll, and sweet potato stem rot, where the first ten-year period shows less loss, and wheat leaf rust, stem rust and crown rust of oats, and peach leaf curl, where the average is the same for both periods.

## WHEAT




Fig. 1. WHEAT: Losses from leaf rust, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.




Fig. 2. WHEAT: Losses from stem rust, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.




Fig. 3. WHEAT: Losses from bunt, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.

## OATS




Fig. 4. OATS: Losses from stem rust, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.




Fig. 5. OATS: Losses from crown rust, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.

OATS - CORN




Fig. 6. OATS: Losses from smut, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.




Fig. 7. CORN: Losses from smut, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.

## CORN




Fig. 8. CORN: Losses from ear rot, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.




Fig. 9. CORN: Losses from root rot, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.

## POTATO

7



Fig. 10. POTATO: Losses from late blight, 1925-34 and averages for the 10-year periods 1920-29 and 1925-34.




Fig. 11. POTATO: Losses from tipburn and hopperburn, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.

## POTATO




Fig. 12. POTATO: Losses from leaf roll, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.




Fig. 13. POTATO: Losses from mosaic, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.




Fig. 14. POTATO: Losses from Rhizoctonia, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.




Fig. 15. SWEET POTATO: Losses from storage rot, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.

## SWEET POTATO





Fig. 17. SWEET POTATO: Losses from black rot, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.

## APPLE




Fig. 18. APPLE: Losses from scab, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.




Fig. 19. APPLE: Losses from bitter rot, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.




Fig. 20. APPLE: Losses from black rot, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.




Fig. 21. APPLE: Losses from blight, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.




Fig. 22. APPLE: Losses from rust, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.




Fig. 23. PEACH: Losses from brown rot, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.




Fig. 24. PEACH: Losses from leaf curl, 1925-34, and averages for the 10-year periods 1920-29 and 1925-34.

## 1934 - WHEAT

## Estimated reduction in yield of wheat due to diseases, 1934

Estimated reduction in yield of wheat due to diseases									
Production Bushels	% Bushels	Scab 1,000	Leaf Rust %	Stem Rust %	Bunt %	Loose Smut %	Foot Rots %	All Diseases Bushels	All Diseases 1,000
Me.	120	-	-	-	-	-	-	-	-
Mass.	4,416	-	+ 10.	+ 10.	-	-	-	-	+ 20.
N. Y.	1,127	-	-	-	-	-	-	-	-
N. J.	14,759	0.5	89	10.	1,783	t	-	-	11
Pa.	33,401	t	+ 0.5	+ 0.5	172	t	-	-	+ 17.2
Ohio	32,152	-	-	-	0	1.5	517	1.	3,067
Ind.	29,495	t	+ 2.	+ 2.	622	0	0	328	1,033
Ill.	11,120	0	0	0	0	0.7	218	1.	656
Mich.	1,647	t	+ 0	+ 0	0	0	2.5	291	1.5
Wis.	12,534	0	0	0	0	0	0.5	678	1.5
Minn.	3,028	0	0	0	0	0	0	68	1.
Iowa	21,281	-	-	-	-	-	-	136	1.
Mo.	21,196	t	+ 2.	+ 2.	448	1.5	-	-	17
N. D.	S. D.	-	-	-	-	-	-	-	-
Nebr.	15,838	-	-	-	-	-	-	-	-
Kans.	79,700	0	0	0	16	t	-	810	0.3
Del.	1,539	1.	1.	1.	448	t	-	269	1.5
Md.	7,934	5.	5.	5.	165	t	-	134	-
Va.	8,092	2.	2.	2.	23	5.	-	70	2.
W. Va.	1,974	1.	1.	1.	107	5.	-	161	1.
N. C.	4,340	2.	2.	2.	143	3.	-	54	3.
S. C.	765	-	-	-	112	5	-	10	1.
Ga.	756	-	-	-	-	-	-	28	3.

Estimated reduction in yield of wheat due to diseases									
Production	1,000 Bushels	Scab	Leaf Rust	Stem Rust	Bunt	Loose Smut	Foot Rots	All Diseases	All Diseases 1,000 Bushels
State	% Bushels	% Bushels	% Bushels	% Bushels	% Bushels	% Bushels	% Bushels	% Bushels	% Bushels
Ky.	4,250	-	-	0	0	-	-	-	-
Tenn.	3,392	t	-	t	-	-	-	4•	141
Ala.	66	-	-	-	-	-	-	-	-
Ark.	297	-	-	-	-	-	-	-	-
Okla.	37,348	-	-	-	-	-	-	-	-
Tex.	25,749	t	+ 5•	1,423	1•	285	1•	2•5	2,705
Mont.	28,174	0	0	t	+	0	0	214	4• 1,221
Idaho	18,696	-	-	-	-	-	-	-	-
Wyo.	1,041	-	-	-	-	-	-	-	-
Colo.	5,776	t	-	0	+	0	0	-	-
N. Mex.	711	-	-	-	-	-	-	-	-
Ariz.	1,000	-	-	-	-	-	-	-	-
Utah	3,147	-	-	-	-	-	-	-	-
Nev.	236	-	-	-	-	-	-	-	-
Wash.	37,246	0	0	t	+	6•	878	t	572
Oreg.	12,544	0	0	0	+	6•	2•	t	293
Calif.	8,284	-	-	-	-	-	-	-	-
U. S.	496,469	0.2	848	1.1	5,517	0.3	1,440	1•	5,382
								0.6	2,999
								0.5	2,491
								4.2	21,125

## Estimated reduction in yield of barley due to diseases, 1934

State	Estimated reduction in yield of barley due to diseases					
	Production:		Stripe	Spot Blotch	Loose Smut	Covered Smut
	Bushels	%	Bushels	%	Bushels	%
Maine	120	-	-	-	-	-
Vt.	108	-	-	-	-	-
Mass.	-	t	+	-	t	+
N.Y.	3,969	-	-	-	-	-
N.J.	27	-	-	-	-	-
Pa.	1,734	1.7	35	1.5	31	3.3
Ohio	332	-	-	-	1.	3
Ind.	273	-	-	-	-	-
Ill.	884	t	+	t	t	+
Mich.	3,384	0	0	t	0.5	17
Wis.	19,266	0.5	101	t	2.5	506
Minn.	24,115	0.5	128	t	1.	255
Iowa	5,712	t	+	1.	58	1.
Mo.	220	-	-	-	-	-
N. D.	7,119	0.2	15	t	0.5	36
S. D.	1,778	-	-	-	-	-
Nebr.	2,700	-	-	-	-	-
Kans.	1,988	t	+	-	0.3	6
Del.	-	-	-	-	-	t
Md.	627	0.5	3	-	1.	7
Va.	882	-	-	-	-	5.
W. Va.	80	1.	1	-	2.	2
N. C.	272	2.	6	1.	3	5.
S. C.	-	t	+	-	-	-
Ga.	-	-	-	-	-	7.
Ky.	210	-	-	-	-	-
Tenn.	323	-	-	-	-	-
Okla.	1,265	-	-	-	-	-
Tex.	2,457	0.1	3	t	1.	26
Mont.	2,214	1.	23	t	1.	23
Idaho	4,288	-	-	-	-	-
Wyo.	774	1.	8	-	2.	16
Colo.	3,024	t	+	-	t	t
N. Mex.	120	-	-	-	-	-
Ariz.	638	-	-	-	-	-
Utah	837	-	-	-	-	-
Nev.	140	-	-	-	-	-
Wash.	1,770	-	-	-	-	-
Oreg.	2,597	-	-	t	-	0.2
Calif.	22,682	-	-	-	-	-
U. S.	118,929	0.4	323	0.1	92	1.3
					1,039	0.7
						558

1934 - Barley (continued)

State	Estimated reduction in yield due to diseases				
	Scab		Stem Rust		Other
	: 1,000 : % Bushels	: 1,000 : % Bushels	: 1,000 : % Bushels	: 1,000 : % Bushels	: 1,000 : % Bushels
Maine	-	-	-	-	-
Vt.	-	-	-	-	-
Mass.	0	0	2.	+	5.
N.Y.	-	-	-	-	-
N.J.	-	-	-	-	-
Pa.	0.5	10	t	3.	62 : 15.6: 321
Ohio	2.	7	0.1	1	0.6
Ind.	-	-	0	0	-
Ill.	1.	9	0	0	t
Mich.	t	+	0.5	17	0.2
Wis.	1.8	364	t	+	-
Minn.	t	+	2.	510	1.
Iowa	0	0	t	+	-
Mo.	-	-	-	-	-
N. D.	0	0	t	+	1.
S. D.	-	-	t	+	-
Nebr.	-	-	-	-	-
Kans.	-	-	-	-	-
Del.	-	-	-	-	-
Md.	3.	21	-	-	0.5
Va.	-	-	-	-	-
W. Va.	-	-	-	2.	2 : 7. :
N. C.	2.	6	t	+	3. : 13.5: 42
S. C.	-	-	-	-	-
Ga.	-	-	-	5.	+
Ky.	-	-	-	-	-
Tenn.	-	-	-	-	-
Okla.	-	-	-	-	-
Tex.	0.1	3	0.1	3	2. : 51 : 4.3: 112
Mont.	0	0	0	0	-
Idaho	-	-	-	-	-
Wyo.	-	-	0	0	-
Colo.	-	-	0	0	-
N. M.	-	-	-	-	-
Ariz.	-	-	-	-	-
Utah	-	-	-	-	-
Nev.	-	-	-	-	-
Wash.	0	0	-	-	-
Oreg.	-	-	t	+	4. : 108 : 4.2: 113
Calif.	-	-	-	-	-
U. S.	0.5	420	0.7	531	0.7
				572	4.4: 3,535

1934 - RYE

## Estimated reduction in yield of rye due to diseases, 1934

Production:		Estimated reduction in yield due to diseases											
State:	tion:									Other		All	
	: 1,000	Smut	Ergot	Leaf Rust	Stem Rust	Diseases				Diseases			
	: Bushels:	: 1,000:	: 1,000:	: 1,000:	: 1,000:	: 1,000:				: 1,000:			
	: : %	: Bush-:	: %	: Bush-:	: %	: Bush-:	: %	: Bush-:	: %	: Bush-:	: %		
	: : els	: : els	: : els	: : els	: : els	: : els	: : els	: : els	: : els	: : els	: : els		
Mass.:	: 0	0	1.	+ 2.	+	4.	+	-	-	7.	+		
N.Y.:	250	-	-	-	-	-	-	-	-	-	-		
N.J.:	342	-	-	-	-	-	-	-	-	-	-		
Pa.:	1,344	t	+	t	+ 6.	91	t	+	5.	76	11.	167	
Ohio:	819	-	-	0.1	1	1.	8	0	0	1.	8	2.1	17
Ind.:	1,495	-	-	-	-	-	-	0	0	-	-	-	
Ill.:	630	3.2	21	t	+ 0.3	2	t	+	1.	7	4.5	30	
Mich.:	1,314	0	0	t	+ t	+	t	+	t	+	0.3	4	
Wis.:	1,768	t	+	1.	18	t	+	t	+	t	+	1.	18
Minn.:	2,474	t	+	t	+ 0	0	0	0	0	t	+	t	+
Iowa:	348	-	-	-	-	-	-	0	0	-	-	-	
Mo.:	120	-	-	-	-	-	-	-	-	-	-	-	
N.D.:	1,030	0	0	0.6	6	0	0	t	+	0.5	5	1.1	11
S.D.:	328	-	-	-	-	-	-	0	0	-	-	-	
Nebr.:	728	-	-	-	-	-	-	0	0	-	-	-	
Kans.:	176	-	-	-	-	-	-	-	-	-	-	-	
Del.:	90	-	-	-	-	t	+	-	-	-	-	t	+
Md.:	300	-	-	-	-	-	-	-	-	-	-	-	
Va.:	550	-	-	-	-	-	-	-	-	-	-	-	
W.Va.:	120	t	+	t	+ t	+	t	+	-	-	-	t	+
N.C.:	495	-	-	-	-	t	+	-	-	5.	26	5.	26
S.C.:	60	-	-	-	-	-	-	-	-	-	-	-	
Ga.:	84	-	-	-	-	1.	1.	-	-	-	-	1.	1
Ky.:	138	-	-	-	-	-	-	-	-	-	-	-	
Tenn.:	82	2.	2	-	-	-	-	-	-	-	-	2.	2
Okla.:	54	-	-	-	-	-	-	-	-	-	-	-	
Tex.:	33	t	+	0	0	1.	+	1.	+	0.5	+	2.5	1
Mont.:	210	-	-	-	-	-	-	0	0	-	-	-	
Idaho:	36	-	-	-	-	-	-	-	-	-	-	-	
Wyo.:	72	-	-	-	-	-	-	0	0	-	-	-	
Colo.:	176	-	-	t	+	-	-	0	0	-	-	t	+
Utah:	15	-	-	-	-	-	-	-	-	-	-	-	
Wash.:	119	0	0	-	-	-	-	-	-	-	-	-	
Oreg.:	240	-	-	t	+ 0.3	+	-	-	0.2	+	0.5	1	
U. S.:	16,040	0.2	23	0.2	25	0.9	102	t	+	1.1	122	2.5	278

## Estimated reduction in yield of oats due to diseases, 1934

State	Production:	Estimated reduction in yield due to diseases					
		Smuts	Stem rust	Crown Rust	All diseases		
		1,000 Bushels	% Bushels	% Bushels	% Bushels	% Bushels	% Bushels
Maine	4,680	-	-	-	-	-	-
N. H.	273	-	-	-	-	-	-
Vt.	1,769	-	-	-	-	-	-
Mass.	160	15.	29	2.	4	t	17.
R. I.	64	-	-	-	-	-	-
Conn.	300	-	-	-	-	-	-
N. Y.	23,408	-	-	-	-	-	-
N. J.	1,452	-	-	-	-	-	-
Pa.	24,915	8.9	2,531	t	+ 0.5	142	12.4
Ohio	25,994	2.	537	0.5	134	0.5	134
Ind.	18,225	-	-	0	0	-	-
Ill.	33,319	4.5	1,813	0.5	201	0.5	201
Mich.	28,717	2.	590	0.5	147	0	0
Wis.	65,352	5.	3,443	0.1	69	t	+ 5.1
Minn.	72,703	2.	1,564	5.	3,909	0	0
Iowa	61,250	4.	2,552	t	+ 0	0	4.
Mo.	13,585	-	-	-	-	-	-
N. D.	8,886	0.5	45	t	+ t	+ 1.5	135
S. D.	3,384	-	-	t	+ -	-	-
Nebr.	8,568	-	-	t	+ -	-	-
Kans.	16,094	2.	328	0	0	0	2.
Del.	132	t	+	-	t	+ t	+
Md.	1,320	5.	69	t	+ t	+ 5.	69
Va.	2,398	3.	74	-	-	-	3.
W. Va.	2,052	5.	114	-	- 2.	46	10.
N. C.	3,519	5.	196	1.	39	1.	39
S. C.	6,596	10.	785	1.	79	5.	393
Ga.	6,384	5.	340	-	1.	68	6.
Fla.	104	-	-	-	-	-	-
Ky.	1,650	-	-	-	-	-	-
Tenn.	1,335	-	-	-	-	-	-
Ala.	2,090	-	-	-	-	-	-
Miss.	651	-	-	-	-	-	-
Ark.	2,046	2.	44	t	+ 5.	110	7.
La.	500	-	-	-	-	-	-
Okla.	20,150	-	-	-	-	-	-
Texas	32,466	3.	1,353	4.	1,804	20.	9,018
Mont.	7,344	5.	387	0	0	0	5.
Idaho	4,488	-	-	-	-	-	-
Wyo.	1,743	2.	36	t	+ -	-	2.
Colo.	2,280	t	+ 0	0	t	+ t	+
N. M.	345	-	-	-	-	-	-
Ariz.	288	-	-	-	-	-	-
Utah	832	-	-	-	-	-	-
Nev.	46	-	-	-	-	-	-
Wash.	6,800	t	+ t	+ -	-	-	+
Oreg.	5,904	0.4	53	10.	1,324	35.	4,633
Calif.	2,254	-	-	-	-	-	-
U. S.	528,815	3.6	16,883	1.7	7,710	3.2	14,784
						10.1	47,090

## Estimated reduction in yield of field corn due to diseases, 1934

		Estimated reduction in yield due to diseases							
	Production 1,000 Bushels	Smut	Root Rots (except Diplodia)	Stalk Rots: (except Diplodia)	Diplodia	Bacterial Wilt	All Diseases		
State	% Bushels	% Bushels	% Bushels	% Bushels	% Bushels	% Bushels	% Bushels		
Maine	656	-	-	-	-	-	-	-	-
N. H.	615	-	-	-	-	-	-	-	-
Vt.	2,814	-	-	-	-	-	-	-	-
Mass.	1,517	2.	32	2.	32	2.	16	5.	80
R. I.	410	-	-	-	-	-	-	-	-
Conn.	2,132	-	-	-	-	-	-	-	-
N. Y.	21,286	-	-	-	-	-	-	-	-
N. J.	7,138	-	-	-	-	-	-	-	-
Pa.	52,896	5.	3,306	t	10.	6,612	5.	3,306	20.
Ohio	92,200	3.	2,927	0.5	488	-	-	2.	1,951
Ind.	94,141	1.	1,029	1.	1,029	4.5	4,630	t	2,058
Ill.	146,760	8.	20,243	4.	10,121	13.5	34,160	15.	37,955
Mich.	33,408	4.	1,411	0.5	176	0.2	71	-	1.5
Wis.	73,904	3.	2,584	0.1	86	1.	861	9.9	8,527
Minn.	76,619	4.	3,368	t	+	5.	4,210	+	86.
Iowa	201,480	7.	16,400	-	-	-	3.	7,028	-
Mo.	26,482	-	-	-	-	-	-	-	4.
N. D.	4,904	2.	102	-	-	0.5	25	1.	9,371
S. D.	12,722	-	-	-	-	-	-	-	-
Nebr.	21,363	-	-	-	-	-	-	-	-
Kans.	10,576	1.	108	1.	108	1.	49	5.	1,000
Del.	4,864	t	+ 1.	+ 1.	1,999	5.	-	-	49
Md.	16,995	t	+ 10.	+ 10.	-	-	-	-	2,999
Va.	35,794	-	-	-	-	-	-	-	-
W. Va.	12,128	3.	413	2.	276	3.	413	3.	138

Estimated reduction in yield due to diseases									
	Production:	Smut:	Root Rots:	Ear Rots:	Stalk Rots:	(except:	Diplodia:	Bacterial:	All Diseases
State	: 1,000 Bushels	: % Bushels	: % Bushels	: 1,000 Bushels	: 1,000 Bushels	: (except Diplodia)	: Diplodia	: Wilt	: Bushels:
N. C.	47,580	1.	547	3.	1,641	3.	547	-	7,111
S. C.	20,760	0.5	116	5.	1,160	5.	1,160	+	2,436
Ga.	39,270	5.	2,206	-	5.	2.	1,324	-	4,854
Fla.	6,390	0.5	43	0.5	7.	600	257	2.	187
Ky.	62,832	-	-	-	-	-	-	-	-
Tenn.	58,894	-	-	-	-	-	-	-	-
Ala.	47,950	-	-	-	-	-	-	-	-
Miss.	40,121	-	-	-	-	-	-	-	-
Ark.	15,398	1.	156	-	-	-	-	-	-
La.	16,248	-	-	-	-	-	-	-	-
Oklahoma	11,644	-	-	-	-	-	-	-	-
Tex.	45,873	1.	494	0.1	49	0.5	247	1.	7,817
Mont.	665	t	-	-	-	-	-	-	3,506
Idaho	1,520	-	-	-	0	0	0	-	7
Wyo.	655	1.	7	-	-	-	-	-	-
Colo.	3,368	1.	34	-	-	-	-	-	-
N. M.	1,088	-	-	-	-	-	-	-	-
Ariz.	490	-	-	-	-	-	-	-	-
Utah	304	-	-	-	-	-	-	-	-
Nev.	40	-	-	-	-	-	-	-	-
Wash.	1,122	t	+	0	2	-	-	-	-
Oreg.	1,852	-	-	0.1	-	-	-	-	-
Calif.	2,850	-	-	-	-	-	-	-	-
U. S.	1,380,718	4.6:55,526	1.4:17,259	5.2:63,982	4.3:52,874	1.5:18,616	0.1: 1,602	: 17.5:213,570	9

## 1934 - SWEET CORN

Estimated reduction in yield of sweet corn due to diseases, 1934

		Estimated reduction in yield due to diseases									
		Bacterial									
State:	Short Tons	Production:		Smut		Wilt		Diplodia		All Diseases	
		Short	Tons	%	Tons	Short	Tons	%	Tons	%	Tons
Maine	39,200	-	-	0	0	-	-	-	t	-	+
N. H.	2,200	-	-	-	-	-	-	-	-	-	-
Vt.	2,800	-	-	-	-	-	-	-	-	-	-
Mass.	5.	+	1.	+	2.	+	8.	+	+	+	+
N. Y.	33,600	4.	1,400	t	+	-	-	-	4.	1,400	
Pa.	9,200	5.	667	20.	2,667	5.	667	31.	4,134		
Ohio	39,900	1.	416	3.	1,247	-	-	-	4.	1,663	
Ind.	50,000	-	-	5.	2,632	-	-	-	5.	2,632	
Ill.	89,200	-	-	5.	4,695	-	-	-	5.	4,695	
Mich.	4,500	1.	47	3.	141	t	+	4.2	197		
Wis.	27,400	5.	1,457	t	+	1.	291	6.	1,748		
Minn.	81,300	5.	4,371	0	0	1.	874	7.	6,119		
Iowa	51,300	-	-	-	-	-	-	-	-	-	
N. D.	2.	+	0	0	0	0	0	3.	+	+	
Nebr.	1,200	-	-	-	-	-	-	-	-	-	
Kans.			t	+	+			t	+	+	
Del.	6,000	t	+	-	-	-	-	1.	60		
Md.	43,600	1.	493	1.5	739	9.	4,434	11.5	5,666		
W. Va.		2.	+	5.	+	3.	+	12.	+	+	
Tenn.	5,300	-	-	-	-	-	-	-	-	-	
Texas		1.	+	3.	+	1.	+	7.1	+	+	
Mont.		t	+	0	0	0	0	1.	+	+	
Colo.		2.	+	-	-	-	-	2.	+	+	
Oreg.		-	-	-	-	-	-	t	+	+	
**Other States	8,900	-	-	-	-	-	-	-	-	-	
U. S.	495,600	2.	8,851	2.7	12,121	1.4	6,266	6.3	28,314		

\*Production figures are of sweet corn for manufacture.

\*\*Other States include: Colorado, Idaho, Kansas, Kentucky, Missouri, Montana, Oklahoma, Oregon, South Dakota, Virginia, Washington, Wyoming.

## 1934 - SWEET POTATO

Estimated reduction in yield due to diseases

State	Production:	Estimated reduction in yield due to diseases					
		All		Diseases		Storage Rots	
		Stem Rot	Black Rot	Stem Rot	Black Rot	Stem Rot	Black Rot
		Bushels	%	Bushels	%	Bushels	%
Conn.		-	-	t	+	t	+
N. J.	2,015	-	-	-	-	-	-
Pa.		-	-	t	+	1.	+
Ind.	440	30.	189	t	+	30.	189
Ill.	560	-	-	-	-	-	-
Iowa.	180	20.	46	2.	5	22.	51
Mo.	561	-	-	-	-	-	-
Kans.	325	7.	25	1.	4	8.	29
Del.	810	1.	8	0.5	4	3.5	29
Md.	750	1.5	12	t	+	7.	56
Va.	3,910	7.	301	1.	43	9.	387
W. Va.		-	-	t	+	t	+
N. C.	8,856	3.	295	2.	197	10.	984
S. C.	4,428	t	+	t	+	t	+
Ga.	7,120	-	-	-	-	-	-
Fla.	1,800	1.	22	5.	110	18.	395
Ky.	1,805	-	-	-	-	-	-
Tenn.	6,180	15.	1,091	t	+	15.	1,091
Ala.	7,614	-	-	-	-	-	-
Miss.	7,526	-	-	-	-	-	-
Ark.	1,620	2.	34	2.	34	4.	68
La.	5,840	1.	60	t	+	2.	120
Oklahoma.	770	-	-	-	-	-	-
Tex.	3,300	0.5	20	8.	324	18.5	749
Colo.		t	+	-	-	t	+
Calif.	990	-	-	-	-	-	-
U. S.	67,400	4.9	2,103	1.7	721	9.7	4,148
						17.	4,237

## Estimated reduction in yield of potato due to diseases

		Estimated reduction in yield due to diseases					
	Production:	Mosaic	Leaf Roll	Late Blight	Rhizoctonia		
State	1,000 Bushels	1,000 % Bushels	1,000 % Bushels	1,000 % Bushels	1,000 % Bushels	1,000 % Bushels	
Maine	56,280	2.	1,324	1.5	993	1.	662
N. H.	1,665	-	-	-	-	-	-
Vt.	2,720	-	-	-	-	-	-
Mass.	2,560	5.	191	5.	191	4.	153
R. I.	540	-	-	-	-	-	-
Conn.	2,625	-	-	-	-	-	-
N. Y.	32,550	2.	910	3.	1,366	3.	1,366
N. J.	9,050	-	-	-	-	-	-
Pa.	34,000	3.	1,407	10.	4,690	2.	938
Ohio	11,445	-	-	5.	728	t	+
Ind.	6,200	-	-	-	-	-	-
Ill.	2,704	-	-	-	-	0	0
Mich.	34,304	0.5	211	1.	421	0	0
Wis.	31,320	t	+	0	0	0	t
Minn.	23,380	5.	1,601	+	+	0	0
Ia.	4,851	-	-	-	-	-	3.
Mo.	1,380	-	-	-	-	-	-
N. D.	5,940	1.	63	1.	63	0	0
S. D.	1,290	-	-	-	-	0	-
Nebr.	3,450	-	-	-	-	0	-
Kans.	1,480	t	+	-	-	0	0
Del.	720	0.5	4	+	+	-	t
Md.	3,267	2.	75	2.	75	1.5	56
Va.	13,433	2.	299	2.	299	0	0
W. Va.	3,120	4	184	5.	229	t	+
N. C.	10,672	2.	248	3.	372	t	+
S. C.	2,625	t	+	0	0	0	3.
Ga.	1,360	3.	52	-	-	0	0
Fla.	3,250	1.	39	+	+	6.	234
Ky.	4,480	-	-	-	-	-	-
Tenn.	4,320	-	-	-	-	0	0
Ala.	3,760	-	-	-	-	-	-
Miss.	1,344	-	-	-	-	-	-
Ark.	2,624	-	-	-	-	0	0
La.	2,948	3.	95	-	-	0	0
Okla.	2,760	-	-	-	-	0	0
Tex.	3,780	0.1	4	t	+	t	+
Mont.	1,610	5.	95	t	+	0	0
Idaho	19,240	-	-	-	-	0	0
Wyo.	1,000	2.	35	-	-	0	0
Colo.	5,700	t	+	t	+	0	0
N. M.	490	-	-	-	-	0	0
Ariz.	100	-	-	-	-	0	0
Utah	1,040	-	-	-	-	0	0
Nev.	210	-	-	-	-	0	0
Wash.	7,290	7.5	591	+	+	t	+
Oreg.	5,720	-	-	-	-	-	-
Calif.	8,610	-	-	-	-	-	-
U. S.	385,287	2.	7,428	2.5	9,427	0.9	3,409
						3.4	12,594

1934 - POTATO (Continued)

State:	Estimated reduction in yield due to diseases										All Diseases	
	Blackleg		Fusarium Wilt		Tipburn and Hopperburn		Early Blight		Scab			
	% Bushels	% Bushels	% Bushels	% Bushels	% Bushels	% Bushels	% Bushels	% Bushels	% Bushels	% Bushels		
Maine:	1.	662	0	0	0	0	-	-	2.	1,324	: 15. : 9,931	
N.H.:	1.	17	-	-	-	-	-	-	-	-	: 1. : 17	
Vt.:	-	-	-	-	-	-	-	-	-	-	-	
Mass.:	t	+	t	+	8.	306	2.	76	1.	38	: 33. : 1,260	
R.I.:	-	-	-	-	-	-	-	-	-	-	-	
Conn.:	-	-	-	-	-	-	-	-	-	-	-	
N.Y.:	t	+	t	+	10.	4,552	t	+	3.	1,366	: 28.5 : 12,974	
N.J.:	-	-	-	-	-	-	-	-	-	-	-	
Pa.:	t	+	2.	938	3.	1,407	t	+	0.5	234	: 27.5 : 12,897	
Ohio:	0.2	29	1.5	218	8.	1,165	2.	291	0.5	73	: 21.4 : 3,115	
Ind.:	-	-	-	-	-	-	-	-	-	-	-	
Ill.:	-	-	-	-	-	-	-	-	-	-	-	
Mich.:	2.	843	1.6	674	5.	2,107	0.5	211	5.	2,107	: 18.6 : 7,838	
Wis.:	0	0	0	0	t	+	0	0	5.	1,666	: 6. : 1,999	
Minn.:	t	+	4.	1,281	8.	2,562	2.	641	2.	641	: 27. : 8,648	
Iowa:	-	-	-	-	-	-	-	-	10.	558	: 13. : 725	
Mo.:	-	-	-	-	-	-	-	-	-	-	-	
N. D.:	1.	63	1.5	94	0.2	13	t	+	t	+	: 5.3 : 334	
S. D.:	-	-	-	-	-	-	-	-	-	-	-	
Nebr.:	-	-	-	-	-	-	-	-	-	-	-	
Kans.:	t	+	t	+	-	-	-	-	1.	16	: 8. : 128	
Del.:	-	-	-	-	-	-	t	+	t	+	: 1. : 8	
Md.:	t	+	2.	75	0.5	19	0.5	19	t	+	: 13. : 488	
Va.:	1.	149	t	+	3.	448	t	+	2.	299	: 10. : 1,494	
W. Va.:	2.	92	3.	138	7.	321	t	+	t	+	: 32. : 1,468	
N.C.:	1.	124	t	+	2.	248	1.	124	-	-	: 14. : 1,736	
S.C.:	t	+	0	0	-	-	1.5	41	-	-	: 4.5 : 123	
Ga.:	8.	140	-	-	5.	87	1.	17	-	-	: 22. : 383	
Fla.:	0.5	19	t	+	0	0	0.5	19	0.5	19	: 16.5 : 642	
Ky.:	-	-	-	-	-	-	-	-	-	-	-	
Tenn.:	-	-	t	+	-	-	25.	1,440	-	-	: 25. : 1,440	
Ala.:	-	-	-	-	-	-	-	-	-	-	-	
Miss.:	-	-	-	-	-	-	-	-	-	-	-	
Ark.:	-	-	-	-	15.	463	-	-	-	-	: 15. : 463	
La.:	t	+	-	-	-	-	0.5	16	-	-	: 6.5 : 206	
Okla.:	-	-	-	-	-	-	-	-	-	-	-	
Tex.:	0	0	1.	44	0.5	22	0.5	22	8.	348	: 13.1 : 571	
Mont.:	t	+	2.	38	t	+	t	+	t	+	: 15. : 285	
Idaho:	-	-	-	-	-	-	-	-	-	-	-	
Wyo.:	t	+	3.	53	-	-	t	+	1.	18	: 43. : 755	
Colo.:	-	-	3.5	215	-	-	t	+	1.	61	: 7. : 429	
N.M.:	-	-	-	-	-	-	-	-	-	-	-	
Ariz.:	-	-	-	-	-	-	-	-	-	-	-	
Utah:	-	-	-	-	-	-	-	-	-	-	-	
Nev.:	-	-	-	-	-	-	-	-	-	-	-	
Wash.:	t	+	t	+	-	-	t	+	t	+	: 7.5 : 591	
Oreg.:	-	-	-	-	-	-	-	-	-	-	-	
Calif.:	-	-	-	-	-	-	-	-	-	-	-	
U. S.:	0.6	2,138	1.0	3,768	3.7	13,720	0.8	2,917	2.3	8,768	: 19. : 70,948	

## 1934 - TOMATOES FOR MANUFACTURE

## Estimated reduction in yield of tomatoes due to diseases, 1934

State	Estimated reduction in yield due to diseases																		
	Production:			Septoria			Fusarium			Blossom-end Rot			Curly-top			All Diseases			
	Shrt	Tons*	Blight	Short	Tons	%	Short	Tons	%	Short	Tons	%	Short	Tons	%	Short	Tons	%	
N. Y.	119,200	t	+ t	+ 1.5	1,912	3.5	4,462	0	0	0	6.5	8,286	-	-	-	-	-	-	
N. J.	122,800	-	-	-	-	-	-	-	-	0	0	-	-	-	-	-	-	-	
Pa.	34,300	5.	2,287	1.	457	3.	1,372	8.	3,659	0	0	25.	11,434	-	-	-	-	-	
Ohio	78,500	1.	924	5.	4,618	1.	924	5.	4,618	0	0	15.	13,855	-	-	-	-	-	
Ind.	315,400	-	-	-	-	2.	6,437	-	-	0	0	2.	6,437	-	-	-	-	-	
Ill.	11,100	1.	148	5.	749	-	-	1.	148	0	0	25.	3,700	-	-	-	-	-	
Mich.	17,300	t	+ t	0	538	t	+ t	0	+	0	0	3.5	628	-	-	-	-	-	
Wis.	7,700	-	-	-	0	1.	+ 0	0	0	0	0	1.	-	-	-	-	-	-	
Iowa	Mo.	1,400	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Del.	51,800	1.	523	t	+ t	+ t	+ t	+ t	+ t	0	0	1.	523	-	-	-	-	-	
Md.	184,100	1.	2,597	0.5	1,298	5.	12,983	t	+ t	0	0	29.1	75,561	-	-	-	-	-	
Va.	43,500	t	+ 7.	3,460	1.	494	-	-	-	0	0	12.	5,931	-	-	-	-	-	
W. Va.	S. C.	2.	+ 2.	+ 2.	+ 2.	+ 2.	+ 2.	+ 2.	+ 2.	0	0	10.	-	-	-	-	-	-	
Ga.		t	+ 5.	+ 5.	+ 5.	+ 5.	+ 5.	+ 5.	+ 5.	0	0	15.	-	-	-	-	-	-	
Fla.		10.	+ 3.	+ 3.	+ 3.	+ 3.	+ 3.	+ 3.	+ 3.	0	0	13.	-	-	-	-	-	-	
Ky.		0	0	2.	2.	-	-	-	-	0	0	16.	-	-	-	-	-	-	
Tenn.		-	-	-	-	-	-	-	-	0	0	0	0	0	0	0	0	0	
Ark.		16,200	-	-	-	-	-	-	-	2.	2.	2.	33.	0	0	0	0	0	
La.		1,600	-	-	-	-	-	-	-	5.	5.	5.	+	0	0	0	0	0	
		t	-	-	-	-	-	-	-	1.	1.	1.	+	0	0	0	0	0	

## 1934 - TOMATOES FOR MANUFACTURE (Continued)

Estimated reduction in yield due to diseases									
Production	Septoria	Fusarium	Blossom-end	Curly-top	All Diseases	Short	Rot	Short	Short
State	Short	Blight	Wilt	Early Blight		Tons*	%	Tons	Tons
	%	Tons	%	Short		%	%	%	%
Tex.	2.	+	3.	*	15.	*	0	0	28.
Colo.	11,000	t	+	t	+	t	+	t	+
Utah	23,000	-	-	2.	773	-	-	-	11,597
Wash.	0	0	-	-	-	-	-	-	30.
Oreg.	0	0	-	-	-	-	-	-	25.
Calif.	291,500	-	-	-	-	-	-	-	25.
*Other									
States	50,600								
U. S.	1,389,600	0.6	6,479	1.9:20,607	2.3:24,122	1.2:12,920	1.1:11,597	14.3:150,878	

\*Short tons 2000 lbs.

\*\*Including Connecticut, Florida, Georgia, Idaho, Kansas, Louisiana, Mississippi, Nebraska, New Mexico, North Carolina, Oklahoma, Oregon, South Carolina, Texas, Washington, West Virginia, Wisconsin.

1934 - TOMATOES FOR MARKET

Estimated reduction in yield of tomatoes due to diseases, 1934

Production:		Septoria		Fusarium		Blight		Wilt		Early Blight		Rot		Blossom-end:		Estimated reduction in yield due to diseases		
State	*Bushels	%	Bushels	%	Bushels	%	Bushels	%	Bushels	%	Bushels	%	Bushels	%	Bushels	%	Bushels	%
Miss.	1,380	-	-	-	-	-	-	-	-	-	-	-	-	-	0	0	-	
Ark.	81	-	-	-	-	-	-	-	-	-	-	-	-	-	0	0	2	
La.	160	t	+t	5.	9	2.	4	1.	2.	2.	2	0	0	0	0	0	26	
Tex.	3,374	2.	94	3.	141	15.	141	15.	0	0	0	0	0	0	0	0	1,313	
Mont.	0	0	t	+	+	0	t	+	t	+	t	+	t	+	t	t	+	
Colo.	238	t	+	t	+	t	+	t	+	t	+	t	+	t	+	t	2	
Utah	114	-	-	2.	3	-	-	-	-	-	-	-	-	-	30.	50	32.	
Wash.	357	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	53	
Oreg.	166	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	119	
Calif.	2,890	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	55	
U. S.	18,080	0.8	120	4.7	699	2.2	324	5.4	805	1.5	224	20.5	224	1.5	224	20.5	3,063	

\*Bushels of 53 pounds.

## 1934 - SNAP BEANS FOR MARKET

## Estimated reduction in yield of snap beans due to diseases, 1934

Estimated reduction in yield due to diseases									
	Production : 1,000 Bushels	Anthracnose % Bushels	Bacterial Blight	Mosaic	Root Rots	Rust	Diseases % Bushels	Other % Bushels	All Diseases % Bushels
Mass.	206	1.0	2.0	2.0	2.0	0.5	0.5	0.5	0.5
N. Y.	1,925	0.5	1.0	4.0	9	t	-	-	4.5
N. J.	-	-	-	-	-	-	-	-	-
Pa.	261	5.0	15	2.0	6	0.5	2	6.0	48
Ohio	52	1.0	+	3.0	+	0.5	+	-	3
Ill.	38	0.0	+	4.0	+	10.0	1	-	15.5
Mich.	0	0.0	0	5.0	+	0.5	+	-	4.5
Wis.	0	0.0	0	t	+	0	0	0	4.5
Minn.	0	t	+	1.0	+	2.0	t	+	5.5
N. D.	8	t	+	t	+	2.0	-	-	2.0
Del.	218	t	+	1.5	3	0.5	1	2.5	4.5
Md.	385	0.0	0	1.0	4	-	2.0	-	3.0
Va.	W. Va.	3.0	3.0	1.0	+	3.0	1.0	-	11.0
N. C.	816	1.0	9	1.0	+	1.0	7.0	1.0	131
S. C.	540	0.0	0	1.0	9	2.0	66	1.0	54
Ga.	250	3.0	9	-	6	1.0	5.0	14	37
Fla.	6,376	1.0	76	5.0	380	t	4.0	76	1,216
Tenn.	77	-	-	-	-	-	-	-	-

## 1934 - SNAP BEANS FOR MARKET (Continued)

31

		Estimated reduction in yield due to disease							
Production:		Bacterial:			Mosaic	Root Rot	Rust	Other	All Diseases
State	1,000 Bushels	Anthracnose	Blight		% Bushels				
State	*Bushels	1,000	1,000		1,000	1,000	1,000	1,000	1,000
Ala.	74	-	-	-	-	-	-	-	-
Miss.	321	-	-	-	-	-	-	-	-
Ark.	24	-	-	-	-	-	-	-	-
La.	585	1.	6.	6.	39	-	2.	13	-
Tex.	395	0	0	5.	22	2.	2.	9	2.
Mont.	0	0	0	t	+ 4.	1.	t	+	1.
Colo.	181	0	0	8.	16	2.	4	2.	4
Wash.	0	0	0	t	+ t	+ t	0	0	15.
Calif.	754	-	-	-	-	-	-	-	-
U. S.	13,486	1.	116	4.1	494	0.4	42	4.	473
									0.9
									102
									3.5
									418
									: 13.9: 1,645

\* Bushels of 30 pounds.

## 1934 - SNAP BEANS FOR MANUFACTURE

Estimated reduction in yield of snap beans due to diseases

		Estimated reduction in yield due to diseases									
State:	Production	Anthrac-		Bacterial		Mosaic		Root Rots		All Diseases	
		nose	Blight	Short	Mosaic	Short	Mosaic	Short	Rots	Short	Short
		Tons	%	Tons	%	Tons	%	Tons	%	Tons	%
Maine:	2,300	-	-	-	-	-	-	-	-	-	-
N. Y.:	12,200	0.5	64	4.	511	t	+	t	+	4.5	575
Pa.:	2,000	5.	118	2.	47	0.5	12	6.	142	15.5	367
Ohio:		1.	+	3.	+	0.5	+	-	-	4.5	+
Ind.:	4,300	-	-	-	-	-	-	-	-	-	-
Ill.:		1.	+	4.	+	10.	+	-	-	17.	+
Mich.:	5,800	0	0	5.	307	t	+	0.5	31	5.5	338
Wis.:	7,800	0	0	t	+	1.	79	0	0	1.	79
Minn.:		t	+	1.	+	2.	+	t	+	4.	+
Kans.:		0	0	0	0	-	-	-	-	-	-
Del.:	1,600	t	+	-	-	-	-	0.5	8	0.5	8
Md.:	9,800	t	+	1.5	154	0.5	51	2.5	257	4.5	462
Va.:		0	0	1.	+	-	-	2.	+	3.	+
W. Va.:		3.	+	3.	+	1.	+	3.	+	11.	+
S. C.:	300	0	0	1.	3	1.	3	5.	17	9.	30
Ga.:		3.	+	-	-	-	-	5.	+	13.	+
Fla.:		1.	+	5.	+	t	+	4.	+	16.	+
Tenn.:	1,400	-	-	-	-	-	-	-	-	-	-
Miss.:	2,100	-	-	-	-	-	-	-	-	-	-
Ark.:	1,800	-	-	-	-	-	-	-	-	-	-
La.:	600	1.	7	6.	40	-	-	2.	13	9.	60
Tex.:		0	0	5.	+	2.	+	2.	+	10.	+
Mont.:		0	0	t	+	4.	+	1.	+	5.	+
Idaho:		0	0	-	-	3.	+	-	-	3.	+
Wyo.:		0	0	3.	+	2.	+	t	+	5.	+
Colo.:	1,000	0	0	8.	91	2.	23	2.	23	12.	137
Utah:	1,400	0	0	-	-	-	-	-	-	-	-
Wash.:	1,700	0	0	t	+	t	+	t	+	15.	300
Oreg.:	4,600	0	0	-	-	-	-	-	-	-	-
Calif.:	2,200	0	0	-	-	-	-	-	-	-	-
*Other:											
States:	4,500										
U. S.:	67,400	0.4	189	2.6	1,153	0.4	168	1.1	491	5.3	2,356

\*Including Alabama, Florida, Georgia, Idaho, Illinois, Iowa, Kansas, Kentucky, Minnesota, Missouri, Montana, Nebraska, New Hampshire, New Jersey, Ohio, Oklahoma, Texas, Vermont, Virginia, West Virginia, Wyoming.

## 1934 - DRY BEANS

Estimated reduction in yield of dry beans due to diseases, 1934

State	Production *1,000 bags	Reduction in yield due to diseases						All Diseases :1,000 Bags	
		Anthrac-		Bacterial:		Mosaic:			
		nose	Blight						
		:1,000: Bags	%	:1,000: Bags	%	:1,000: Bags	%	:1,000: Bags	
Maine	62	-	-	-	-	-	-	-	
Vt.	18	-	-	-	-	-	-	-	
N.Y.	891	0.5	5	4.	37	t	+	4.5	
Mich.	3,244	0	0	5.	172	t	0.5	5.5	
Wis.	23	0	0	t	+	1.	+	1.	
Minn.	21	t	+	1.	+	2.	+	4.	
Nebr.	68	-	-	-	-	-	-	-	
Mont.	174	0	0	t	+	4.	1.	2.	
Idaho	1,342	0	0	-	-	3.	42	3.	
Wyo.	250	0	0	1.	3	2.	5	3.	
Colo.	279	0	0	2.5	8	t	+	8	
N. M.	66	0	0	-	-	-	-	-	
Ariz.	40	0	0	-	-	-	-	-	
Oreg.	6	0	0	-	-	-	-	-	
Calif.	3,675	0	0	-	-	-	-	-	
U. S.	10,159	t	5	3.4	220	0.8	54	0.5	
							34	4.8	
								314	

\*Bags of 100 pounds.

## 1934 - GREEN PEAS FOR MARKET

Estimated reduction in yield of green peas for market

State	Production Bushels:	Estimated reduction in yield due to disease										
		Bacterial Blight			Fusarium Wilt			Ascochyta Blight			Root Rots	
		%	Bush- els	%	Bush- els	%	Bush- els	%	Bush- els	%	All Diseases	
Mass.		1.	+	1.	+	1.	+	25.	+	30.	+	
N. Y.	456	t	+	t	+	t	+	0.5	2	0.5	2	
N. J.	200	-	-	-	-	-	-	-	-	-	-	
N. D.		-	-	t	+	0	0	t	+	t	+	
Md.	14	0.2	+	0.5	+	0.2	+	3.5	+	4.4	+	
Va.	72	-	-	-	-	-	-	-	-	-	-	
N. C.	228	1.	3	2.	5	2	5	2.	5	3.	23	
S. C.	120	-	-	-	-	-	-	25.	40	25.	40	
Ga.		10.	+	-	-	-	10	15.	+	35.	+	
Fla.	461	t	+	1.	5	2	10	1.	5	8.	40	
Ala.	10	-	-	-	-	-	-	-	-	-	-	
Miss.	138	-	-	-	-	-	-	-	-	-	-	
La.	64	-	-	-	-	-	-	-	-	-	-	
Texas	65	t	+	0	0	0	0	2.	1	3.	2	
Idaho	170	-	-	1.	2	-	-	2.	4	3.	6	
Colo.	673	3.	22	-	-	3.	22	2.	15	8.	59	
Ariz.	126	-	-	-	-	-	-	-	-	-	-	
Wash.	798	t	+	t	+	-	-	t	+	t	+	
Oreg.	48	-	-	-	-	-	-	-	-	-	-	
Calif.	3,849	-	-	-	-	-	-	-	-	-	-	
U. S.	7,442	0.8	25	0.4	12	1.2	37	2.3	72	5.5	172	

\*Bushels of 30 pounds.

## 1934 - GREEN PEAS FOR MANUFACTURE

Estimated reduction in yield of green peas for manufacture

State	Estimated reduction in yield due to disease											
	Production		Bacterial Blight	Fusarium Wilt	Ascochyta Blight		Root Rots		All Diseases			
	Short	Tons	Short	Tons	Short	Tons	Short	Tons	Short	Tons	Short	Tons
	:	:	:	:	:	:	:	:	:	:	:	:
Maine	2,330	-	-	-	-	-	-	-	-	-	-	-
N. Y.	13,280	t	+	t	+	t	+	0.5	67	0.5	67	
Pa.	3,000	-	-	t	+	t	+	20.	750	20.	750	
Ohio	1,540	-	-	-	-	-	-	1.	16	1.	16	
Ind.	2,610	-	-	-	-	-	-	-	-	-	-	
Ill.	2,070	l.	22	l.	22	-	-	5.	111	7.	155	
Mich.	6,390	0	0	t	+	t	+	1.	65	1.	65	
Wis.	71,120	t	+	0.5	359	0	0	0.5	359	1.	718	
Minn.	5,800	-	-	-	-	-	-	-	-	-	-	
Del.	3,010	-	-	-	-	-	-	-	-	-	-	
Md.	14,850	0.2	31	0.5	78	0.2	31	3.5	544	4.4	684	
Mont.	2,620	1.	32	0	0	t	+	16.	505	17.	537	
Idaho		-	-	1.	+	-	-	2.	+	3.	+	
Wyo.		t	+	-	-	t	+	2.	+	2.	+	
Colo.	3,430	3.	112	-	-	3.	112	2.	75	8.	299	
Utah	11,020	-	-	-	-	-	-	-	-	-	-	
Wash.	9,030	t	+	t	+	-	-	t	+	t	+	
*Other States	12,670											
U. S.	164,770	0.1	197	0.4	459	0.1	143	1.9	2,492	2.5	3,291	

\*Including California, Idaho, Iowa, Kansas, Nebraska, New Jersey, Oklahoma, Oregon, Tennessee, Virginia, Wyoming.

### Estimated reduction in yield of cotton due to diseases, 1934

## 1934 - SUGAR BEET

Estimated reduction in yield of sugar beet due to diseases, 1934

Estimated reduction in yield due to diseases											
Production		Nematode	Curly-top	Leaf Spot	Phoma Rot	Diseases	Diseases	Other	All		
State:	Short	: 1,000:	: 1,000:	: 1,000:	: 1,000:	: 1,000:	: 1,000:	: 1,000:	: 1,000:		
	Tons	%	Short	%	Short	%	Short	%	Short	%	Short
Ohio	329	0	0	0	0	1.	3	1.	3	2.	6
Mich.	1,081	0	0	0	0	t	+	2.	22	t	2.1
Wis.	162	0	0	0	0	t	+	t	+	t	+
Minn.		0	0	0	0	1.	+	t	+	t	+
Nebr.	550	-	-	-	-	-	-	-	-	-	-
Mont.	768	-	-	t	+	t	+	1.	8	1.	8
Idaho	298	-	-	-	-	-	-	-	-	-	-
Wyo.	434	t	+	-	-	-	-	-	t	+	-
Colo.	1,569	-	-	-	-	-	-	-	-	-	-
Utah	261	-	-	-	-	-	-	-	-	-	-
Wash.		-	-	15.	+	0	0	-	-	-	15.
Oreg.		-	-	75.	+	-	-	-	-	-	75.
Calif.	1,596	-	-	-	-	-	-	-	-	-	-
Other:											
States:	579										
*U. S.:	7,627	:	:	:	:	:	:	:	:	:	:

\*Not enough information to calculate losses for the country.

## 1934 - APPLE

Estimated reduction in yield of apples due to diseases

		Estimated reduction in yield due to diseases					
State	Production:			Black Rot			Blotch
	1,000	Bitter Rot	1,000	1,000	1,000	1,000	
	Bushels	%	Bushels	%	Bushels	%	Bushels
Maine	556	-	-	-	-	-	-
N. H.	312	-	-	-	-	-	-
Vt.	255	-	-	-	-	-	-
Mass.	1,435	1.	16	3.	49	0	0
R. I.	97	-	-	-	-	-	-
Conn.	356	-	-	-	-	-	-
N. Y.	11,844	-	-	-	-	-	-
N. J.	2,070	-	-	-	-	-	-
Pa.	8,554	0.1	11	2.	213	0.2	21
Ohio	4,032	2.	84	0.2	8	0.1	4
Ind.	1,020	0.1	1	t	+	0.1	1
Ill.	2,162	0	0	0.5	11	0.5	11
Mich.	6,464	0	0	t	+	0	0
Wis.	1,204	0	0	t	+	t	+
Minn.	339	0	0	0	0	0	0
Iowa	962	-	-	-	-	t	+
Mo.	1,534	-	-	-	-	-	-
N. D.	-	0	0	0	0	0	0
S. D.	34	-	-	-	-	-	-
Nebr.	209	-	-	-	-	-	-
Kans.	783	-	-	t	+	t	+
Del.	688	t	+	0.5	4	t	+
Md.	1,102	2.	26	t	+	0.5	6
Va.	9,275	1.	112	t	+	t	+
W. Va.	3,630	t	+	t	+	t	+
N. C.	3,525	2.	92	t	+	2.	92
S. C.	244	3.	8	0.5	1	-	-
Ga.	874	5.	57	3.	34	-	-
Ky.	934	-	-	-	-	-	-
Tenn.	1,748	-	-	2.5	52	t	+
Ala.	588	-	-	-	-	-	-
Miss.	133	-	-	-	-	-	-
Ark.	1,715	0	0	-	-	0	0
La.	12	-	-	-	-	-	-
Okla.	343	-	-	-	-	-	-
Tex.	134	t	+	t	+	t	+
Mont.	358	0	0	0	0	0	0
Idaho	3,312	0	0	0	0	0	0
Wyo.	28	0	0	0	0	0	0
Colo.	1,354	0	0	0	0	0	0
N. M.	1,240	0	0	0	0	0	0
Ariz.	63	0	0	0	0	0	0
Utah	554	0	0	0	0	0	0
Nev.	36	0	0	0	0	0	0
Wash.	32,300	0	0	-	-	0	0
Oreg.	4,938	0	0	-	-	0	0
Calif.	6,500	0	0	-	-	0	0
U. S.	119,855	0.4	407	0.4	372	0.1	135

1934 - APPLE (Continued)

State	Cedar Rust		Blight		Scab		All Diseases	
	1,000		1,000		1,000		1,000	
	%	Bushels	%	Bushels	%	Bushels	%	Bushels
Maine	-	-	-	-	-	-	-	-
N. H.	-	-	-	-	-	-	-	-
Vt.	-	-	-	-	-	-	-	-
Mass.	3.	49	2.	33	3.	49	12.5	204
R. I.	-	-	-	-	-	-	-	-
Conn.	-	-	-	-	-	-	-	-
N. Y.	-	-	-	-	-	-	-	-
N. J.	-	-	-	-	-	-	-	-
Pa.	0.5	53	1.	107	15.	1,600	19.8	2,112
Ohio	0.1	4	0.5	21	1.	42	4.1	171
Ind.	-	-	2.	21	0.	0	2.2	23
Ill.	t	+	0.5	11	1.	22	2.5	55
Mich.	0	0	t	+	1.	66	2.	132
Wis.	t	+	2.	25	1.	12	3.	37
Minn.	t	+	t	+	t	+	t	+
Iowa	-	-	-	-	0	0	t	+
Mo.	-	-	-	-	-	-	-	-
N. D.	t	+	1.5	+	t	+	1.5	+
S. D.	-	-	-	-	-	-	-	-
Nebr.	-	-	-	-	-	-	-	-
Kans.	-	-	t	+	t	+	t	+
Del.	t	+	t	+	4.	29	5.5	40
Md.	0.5	6	3.	38	5.	64	14.	178
Va.	2.	224	1.	112	8.	894	17.	1,901
W. Va.	1.	38	t	+	2.	76	5.	190
N. C.	1.	46	10.	458	3.	137	23.	1,054
S. C.	-	-	2.	5	t	+	7.	18
Ga.	-	-	10.	114	5.	57	23.	262
Ky.	-	-	-	-	-	-	-	-
Tenn.	-	-	5.	105	5.	105	16.5	346
Ala.	-	-	-	-	-	-	-	-
Miss.	-	-	-	-	-	-	-	-
Ark.	-	-	-	-	t	+	t	+
La.	-	-	-	-	-	-	-	-
Okla.	-	-	-	-	-	-	-	-
Tex.	t	+	2.	3	0	0	7.	10
Mont.	0	0	1.	4	1.	4	2.	8
Idaho	0	0	-	-	-	-	-	-
Wyo.	0	0	-	-	-	-	-	-
Colo.	0	0	t	+	0	0	t	+
N. M.	0	0	-	-	0	0	-	-
Ariz.	0	0	-	-	0	0	-	-
Utah	0	0	-	-	0	0	-	-
Nev.	0	0	-	-	0	0	-	-
Wash.	0	0	t	+	1.	333	3.	999
Oreg.	0	0	t	+	2.	106	7.	372
Calif.	0	0	-	-	-	-	-	-
U. S.	0.4	420	1.1	1,057	3.7	3,596	8.3	8,112

### Estimated reduction in yield of pear due to diseases

State	Estimated reduction in yield due to diseases					
	Production:		Blight	Scab	Leaf Blight	All Diseases
	1,000 Bushels	% Bushels	1,000 Bushels	% Bushels	1,000 Bushels	% Bushels
Maine	2	t	+	-	-	-
N. H.	5	-	-	-	-	-
Vt.	3	-	-	-	-	-
Mass.	35	7	3	1	+*	-
R. I.	6	-	-	-	-	-
Conn.	14	-	-	-	-	-
N. Y.	1,011	-	-	-	-	-
N. J.	74	-	-	-	-	-
Pa.	385	12	56	1	5	14
Ohio	394	0.5	2	0.1	+*	-
Ind.	178	-	-	-	-	-
Ill.	641	5	34	t	+*	5
Mich.	745	1	3	t	+*	1
Wis.	-	1	+*	2	+*	3
Iowa	74	-	-	-	-	-
Mo.	298	-	-	-	-	-
Nebr.	20	-	-	-	-	-
Kans.	145	t	+*	-	-	t
Del.	17	t	+*	-	-	1
Md.	94	1	1	0.5	+*	4
Va.	194	5	10	t	+*	6
W. Va.	25	8	2	2	1	10
N. C.	218	15	40	-	-	3
S. C.	84	-	-	-	-	-
Ga.	185	-	-	-	-	-
Fla.	67	-	-	-	-	-
Ky.	184	-	-	-	-	-
Tenn.	285	20	71	-	-	20
Ala.	326	-	-	-	-	-
Miss.	323	-	-	-	-	-
Ark.	158	-	-	-	-	-
La.	69	5	4	-	-	10
Okla.	171	-	-	-	-	-
Texas	303	4	13	0	0	9
Idaho	52	-	-	-	-	-
Colo.	307	t	+*	0	0	t
N. M.	49	-	-	0	0	-
Ariz.	11	-	-	0	0	-
Utah	53	-	-	0	0	-
Nev.	6	-	-	0	0	-
Wash.	4,042	t	+*	1	41	1
Oreg.	2,470	3	81	2	54	8
Calif.	9,751	-	-	-	-	216
U. S.	23,474	3.0	325	0.9	101	0.2
					22	5.2
						565

## Estimated reduction in yield of peach due to diseases

State	Production	Estimated reduction in yield due to diseases												
		Leaf Curl			Brown Rot			Diseases			Scab			All Diseases
		Bushels	%	Bushels	%	Bushels	%	Bushels	%	Bushels	%	Bushels	%	Bushels
Maine	-	t	+	t	+	-	-	-	-	-	-	t	+	
Mass.	2	2.	+	t	+	t	+	t	+	t	+	2.	+	
R. I.	2	-	-	-	-	-	-	-	-	-	-	-	-	
Conn.	2	-	-	-	-	-	-	-	-	-	-	-	-	
N. Y.	41	-	-	-	-	-	-	-	-	-	-	-	-	
N. J.	22	-	-	-	-	-	-	-	-	-	-	-	-	
Pa.	442	8.	48	15.	90	1.	6	1.	6	26.	156			
Ohio	228	t	+	1.	2	-	-	1.	2	2.1	4			
Ind.	192	t	+	-	-	-	-	t	+	-	-			
Ill.	528	0.5	3	1.	5	t	+	t	+	1.5	8			
Mich.	423	0	0	t	+	0.5	2	t	+	0.5	2			
Iowa	53	-	-	-	-	-	-	-	-	-	-			
Mo.	468	-	-	-	-	-	-	-	-	-	-			
Nebr.	5	-	-	-	-	-	-	-	-	-	-			
Kans.	90	-	-	-	-	-	-	-	-	-	-			
Del.	64	t	+	5.	3	t	+	1.	1	6.	4			
Md.	82	t	+	0.5	+	t	+	0.5	+	1.	1			
Va.	414	t	+	t	+	t	+	-	-	2.	8			
W. Va.	110	t	+	1.	1	-	-	t	+	2.	2			
N. C.	2,312	0.5	14	5.	137	-	-	3.	82	15.5	425			
S. C.	1,610	t	+	1.5	26	-	-	5.	86	6.5	112			
Ga.	5,610	1.	61	2.	122	-	-	5.	305	8.	488			
Fla.	68	-	-	-	-	-	-	-	-	-	-			
Ky.	434	-	-	-	-	-	-	-	-	-	-			
Tenn.	2,325	-	-	10.	258	-	-	0	0	10.	258			
Ala.	1,089	-	-	-	-	-	-	-	-	-	-			
Miss.	781	-	-	-	-	-	-	-	-	-	-			
Ark.	1,848	2.	38	0.5	9	-	-	0	0	2.5	47			
La.	198	-	-	-	-	-	-	-	-	-	-			
Okla.	612	-	-	-	-	-	-	-	-	-	-			
Tex.	1,287	t	+	0.5	7	t	+	3.	42	8.5	119			
Idaho	93	-	-	0	0	-	-	0	0	-	-			
Colo.	1,260	0	0	t	+	t	+	0	0	t	+			
N. M.	123	-	-	0	0	-	-	0	0	-	-			
Ariz.	48	-	-	0	0	-	-	0	0	-	-			
Utah	558	-	-	0	0	-	-	0	0	-	-			
Nev.	6	-	-	0	0	-	-	0	0	-	-			
Wash.	1,200	1.	12	1.	12	-	-	0	0	2.	24			
Oreg.	314	3.	10	2.	7	-	-	0	0	7.	24			
Calif.	20,460	-	-	-	-	-	-	-	-	-	-			
U. S.	45,404	0.9	186	3.1	679	t	8	2.4	524	7.7	1,682			

## 1934 - CHERRY

Estimated reduction in yield of cherry due to diseases

State	Production Tons	Estimated reduction in yield due to diseases									
		Production:		Brown Rot		Leaf Spot		Other Diseases		All Diseases	
		%	Tons	%	Tons	%	Tons	%	Tons	%	Tons
Mass.		20.	+	t	+	-	-	-	20.	+	
N. Y.	19,220	-	-	-	-	-	-	-	-	-	
Pa.	6,344	10.	813	10.	813	2.	163	22.	1,789		
Ohio	3,660	t	+	t	+	-	-	t		+	
Ind.		t	+	t	+	-	-	t		+	
Ill.		2.	+	t	+	-	-	2.		+	
Mich.	26,560	t	+	1.	268	t	+	1.	268		
Wis.	4,400	t	+	3.	136	-	-	3.	136		
Kans.		t	+	-	-	-	-	t		+	
Del.		5.	+	10.	+	-	-	15.		+	
Md.		2.	+	1.	+	-	-	3.		+	
Va.		1.	+	2.	+	-	-	3.		+	
W. Va.		2.	+	t	+	t	+	3.		+	
N. C.		20.	+	t	+	2.	+	22.		+	
Mont.	750	0	0	t	+	-	-	t		+	
Idaho	2,982	0	0	-	-	-	-	-		-	
Colo.	5,920	t	+	-	-	-	-	-		-	
Utah	3,850	-	-	-	-	-	-	-		-	
Wash.	13,500	5.	711	t	+	-	-	5.	711		
Oreg.	9,620	7.	811	5.	580	5.	580	17.	1,971		
Calif.	16,000	-	-	-	-	-	-	-		-	
Twelve:											
States:	112,806	3.3	2,335	2.6	1,797	1.1	743	7.0	4,875		

## 1934 - GRAPE

Estimated reduction in yield of grape due to diseases  
east of Rocky Mountains

State	Production:	Estimated reduction in yield due to diseases					
		Black Rot		Other Diseases		All Diseases	
		Tons	%	Tons	%	Tons	%
Maine	7	-	-	-	-	-	-
N. H.	29	-	-	-	-	-	-
Vt.	11	-	-	-	-	-	-
Mass.	307	8.	27	t	+	8.	27
R. I.	171	-	-	-	-	-	-
Conn.	1,023	-	-	-	-	-	-
N. Y.	49,400	-	-	-	-	-	-
N. J.	2,464	-	-	-	-	-	-
Pa.	18,981	12.	2,649	2.	441	14.	3,090
Ohio	22,720	2.	464	t	+	2.	464
Ind.	2,812	-	-	-	-	-	-
Ill.	5,658	1.	57	-	-	1.	57
Mich.	61,145	t	+	1.	618	1.	618
Wis.	274	1.	3	-	-	1.	3
Minn.	194	-	-	t	+	t	+
Iowa	5,060	-	-	-	-	-	-
Mo.	7,540	-	-	-	-	-	-
Nebr.	1,216	-	-	-	-	-	-
Kans.	2,574	t	+	-	-	t	+
Del.	2,430	20.	608	-	-	20.	608
Md.	614	15.	108	t	+	15.	108
Va.	1,692	15.	317	5.	106	20.	423
W. Va.	944	5.	52	5.	52	10.	104
N. C.	4,640	10.	546	5.	273	15.	819
S. C.	829	25.	276	-	-	25.	276
Ga.	738	15.	148	10.	98	25.	246
Fla.	1,026	10.	128	10.	128	20.	256
Ky.	1,113	-	-	-	-	-	-
Tenn.	1,110	50.	1,110	-	-	50.	1,110
Ala.	641	-	-	-	-	-	-
Miss.	228	-	-	-	-	-	-
Ark.	16,640	t	+	-	-	t	+
La.	44	-	-	-	-	-	-
Okla.	2,112	-	-	-	-	-	-
Texas	1,595	5.	92	8.	147	13.	239
	217,982	4.3	6,585	1.2	1,863	5.5	8,448

## 1934 - STRAWBERRY

## Estimated reduction in yield due to diseases, 1934

State	Production: 1,000 *Crates	Estimated reduction in yield due to diseases										Fruit Rots: % : Crates	
		Leaf Spot:			Dwarf Scorch:			Black Root:			All Diseases:		
		1,000 *Crates	%	Crates:	1,000 *Crates	%	Crates:	1,000 *Crates	%	Crates:	1,000 *Crates	%	
Maine	1.	+	-	0	0	-	-	-	-	1.	+	-	-
Mass.	2.	+	-	t	+	9.	+	12.	+	23.	+	-	+
N. Y.	-	-	-	-	-	-	-	-	-	-	-	-	-
N. J.	-	-	-	-	-	-	-	-	-	-	-	-	-
Pa.	260	2.	6	-	-	0	8.	-	-	-	-	-	-
Ohio	189	0.1	+	0	0	0	-	23	1.	3	11.	32	-
Ind.	74	-	-	-	-	-	-	-	-	-	-	-	-
Ill.	320	t	-	-	-	-	-	-	-	-	-	-	-
Mich.	285	t	+	t	+	0	0	5.	t	1.	25.	58	-
Wis.	173	t	+	t	+	0	0	0	1.	3	1.	1.	-
Minn.	66	2.	0	0	0	0	0	25.	1.	2.	5.	-	-
Iowa	720	-	-	-	-	-	-	-	-	-	-	-	-
Mo.	N. D.	16	t	+	-	0	0	-	-	1.	1.	-	-
Kans.	352	-	-	-	-	0	0	-	-	-	-	-	-
Del.	652	1.	8	1.	8	-	-	15.	1.	17.	17.	134	-
Md.	732	5.	44	2.	18	-	-	10.	1.	88	3.	16.	150
Va.	N. C.	476	t	+ 1.	6	2.	11	: 10.	57	3.	17	91	8.
S. C.	Ga.	39	-	-	-	-	-	-	-	-	-	-	-
Fla.		27	-	-	-	-	-	-	-	-	-	-	-
		675	0	0	t	1.	8	1.5	12	12.5	99	15.	119

1934 - STRAWBERRY (Continued)

State	Production: 1,000 *Crates	Estimated reduction in yield due to diseases										All Diseases % : Crates: 1,000: % : Crates:	
		Leaf Spot			Leaf Scorch			Dwarf			Black Root and Root Rots:		
		% : Crates:	1,000:	% : Crates:	% : Crates:	1,000:	% : Crates:	% : Crates:	1,000:	% : Crates:	1,000:	% : Crates:	
Ky.	533	-	-	-	-	-	-	-	-	-	-	-	-
Tenn.	1,048	-	-	-	-	-	-	-	-	-	-	-	-
Ala.	220	-	-	-	-	-	-	-	-	-	-	-	-
Miss.	77	-	-	-	-	-	-	-	-	-	-	-	-
Ark.	1,518	-	-	-	-	-	-	-	-	-	-	-	-
La.	1,242	10.	148	5.	74	0.5	7	0.5	7	-	-	16.	186
Okla.	76	-	-	-	-	-	-	-	-	-	-	-	-
Texas	184	1.	2	0.5	1	0	0	0	3.	6	5.	10	9.5
Mont.	-	t	+	0	0	0	0	0	5.	+	11.	+	-
Colo.	-	-	-	-	-	-	-	-	0	t	-	t	-
Utah	43	-	-	-	-	-	-	-	-	-	-	-	-
Wash.	638	-	-	0	0	0	0	0	7.5	55	5.	36	12.5
Oreg.	595	2.	15	0	0	0	0	0	5.	38	15.	114	22.
Calif.	989	-	-	-	-	-	-	-	-	-	-	-	-
U. S. :	13,264	3.0:	223	1.4:	107	0.3:	26	6.4:	482	3.7:	279	14.8:	1,117
													224

\*24-quart crates, 36 pounds.

